

Beaune, D., & Sellier, Y. (2021a). Stream restorations with meanders increase dragonfly and damselfly diversity and abundance, including an endangered species. *Journal for Nature Conservation*, 60. <https://doi.org/10.1016/j.jnc.2020.125950>

Beaune, D., & Sellier, Y. (2021b). Stream restorations with meanders increase dragonfly and damselfly diversity and abundance, including an endangered species. *Journal for Nature Conservation*, 60. <https://doi.org/10.1016/j.jnc.2020.125950>

Boudot, J.-P., Cugno, A., Dommanget, J.-L., Faton, J.-M., Ga-Zay, C., Mouquet, C., Olivereau, F., Papazian, M., Piney, B., Rusu-Stie-Venard, A., & Sobczyk-Moran, G. (2020).

Plan national d'actions 2020-2030 En faveur des libellules.

Bower, L. M., Stoczynski, L., Peoples, B. K., Patrick, C. J., & Brown, B. L. (2023a). Multiple dimensions of functional diversity affect stream fish taxonomic β -diversity. *Freshwater Biology*, 68(3), 437–451. <https://doi.org/10.1111/fwb.14036>

Bower, L. M., Stoczynski, L., Peoples, B. K., Patrick, C. J., & Brown, B. L. (2023b). Multiple dimensions of functional diversity affect stream fish taxonomic β -diversity. *Freshwater Biology*, 68(3), 437–451. <https://doi.org/10.1111/fwb.14036>

Bretagne Vivante. (2017). *Atlas de répartition provisoire des odonates de Bretagne.*

Cerema. (2019). *Infrastructures linéaires de transport et Odonates.*

Csutoros, A. (2023). *Première mention de Leucorrhinia pectoralis (Charpentier, 1825) en Bretagne (Odonata : Libellulidae).*

Cunningham-Minnick, M. J., Meyer, T. B., & Crist, T. O. (2019a). Shifts in dragonfly community structure across aquatic ecotones. *International Journal of Odonatology*, 22(2), 121–133. <https://doi.org/10.1080/13887890.2019.1615006>

Cunningham-Minnick, M. J., Meyer, T. B., & Crist, T. O. (2019b). Shifts in dragonfly community structure across aquatic ecotones. *International Journal of Odonatology*, 22(2), 121–133. <https://doi.org/10.1080/13887890.2019.1615006>

Érienne Orio, C. M. (n.d.). *Entomofaune des cours d'eau et des zones humides : les Odonates.*
GRETIA. (2010). *Synthèse des connaissances préalable à la Déclinaison régionale du Plan national d'actions Odonates en Basse-Normandie.* www.insectes.org

GRETIA. (2015a). *Bien gérer ses rivières pour la Cordulie à corps fin (Oxygastra curtisii) en Basse-Normandie.* www.basse-normandie.developpement-durable.gouv.fr

GRETIA. (2015b). *Éléments de doctrine régionale pour la prise en compte des odonates dans le cadre des études réglementaires en Pays de la Loire.*

Hernández, M. Á., Casanueva, P., Nunes, L., Santamaría, T., Sánchez-Sastre, L. F., Ferreras-Romero, M., & Campos, F. (2022a). Geographical variation of prementum size in Iberian *Cordulegaster boltonii* (Odonata: Cordulegastridae) populations. *International Journal of Odonatology*, 25, 56–61. <https://doi.org/10.48156/1388.2022.1917169>

Hernández, M. Á., Casanueva, P., Nunes, L., Santamaría, T., Sánchez-Sastre, L. F., Ferreras-Romero, M., & Campos, F. (2022b). Geographical variation of prementum size in Iberian *Cordulegaster boltonii* (Odonata: Cordulegastridae) populations. *International Journal of Odonatology*, 25, 56–61. <https://doi.org/10.48156/1388.2022.1917169>

Kietzka, G. J., Pryke, J. S., & Samways, M. J. (2018a). Comparative effects of urban and agricultural land transformation on Odonata assemblages in a biodiversity hotspot. *Basic and Applied Ecology*, 33, 89–98. <https://doi.org/10.1016/j.baae.2018.08.008>

Kietzka, G. J., Pryke, J. S., & Samways, M. J. (2018b). Comparative effects of urban and agricultural land transformation on Odonata assemblages in a biodiversity hotspot. *Basic and Applied Ecology*, 33, 89–98. <https://doi.org/10.1016/j.baae.2018.08.008>

Leipelt, K. G., & Suhling, F. (2001). *Habitat selection of larval Gomphus gralinii and Oxygastra curtisii* (Odonata: Gomphidae, Corduliidae).

Mondy, C. P., Villeneuve, B., Archaimbault, V., & Usseglio-Polatera, P. (2012a). A new macroinvertebrate-based multimetric index (I 2M 2) to evaluate ecological quality of French wadeable streams fulfilling the WFD demands: A taxonomical and trait approach. *Ecological Indicators*, 18, 452–467. <https://doi.org/10.1016/j.ecolind.2011.12.013>

Office National des Forêts, A. de l'Eau S. N. A. de l'Eau A. P. C. R. d'Ile-de-F. (2006). *La gestion des mares forestières de plaine*.

OPIE. (2016). *Aborder la gestion conservatoire en faveur des Odonates*. www.odonates.pnaopie.fr

Picard, L. (2020). *A propos des invertébrés de Bretagne*.

Valente-Neto, F., Rodrigues, M. E., & Roque, F. de O. (2018a). Selecting indicators based on biodiversity surrogacy and environmental response in a riverine network: Bringing operationality to biomonitoring. *Ecological Indicators*, 94, 198–206. <https://doi.org/10.1016/j.ecolind.2018.06.066>

Valente-Neto, F., Rodrigues, M. E., & Roque, F. de O. (2018b). Selecting indicators based on biodiversity surrogacy and environmental response in a riverine network: Bringing operationality to biomonitoring. *Ecological Indicators*, 94, 198–206. <https://doi.org/10.1016/j.ecolind.2018.06.066>

Vilenica, M., Pozojević, I., Vučković, N., & Mihaljević, Z. (2020a). How suitable are man-made water bodies as habitats for Odonata? *Knowledge and Management of Aquatic Ecosystems*, 2019-January(421). <https://doi.org/10.1051/kmae/2020008>

Vilenica, M., Pozojević, I., Vučković, N., & Mihaljević, Z. (2020b). How suitable are man-made water bodies as habitats for Odonata? *Knowledge and Management of Aquatic Ecosystems*, 2019-January(421). <https://doi.org/10.1051/kmae/2020008>